

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 403/06, 413/14, A61K 31/42, 31/405		A1	(11) International Publication Number: WO 96/17842 (43) International Publication Date: 13 June 1996 (13.06.96)
(21) International Application Number: PCT/GB95/02759 (22) International Filing Date: 27 November 1995 (27.11.95) (30) Priority Data: 9424627.9 6 December 1994 (06.12.94) GB 9505871.5 23 March 1995 (23.03.95) GB (71) Applicant (for all designated States except US): MERCK SHARP & DOHME LIMITED (GB/GB); Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): CASTRO PINEIRO, Jose, Luis (ES/GB); Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). HENRY, Caroline (GB/GB); 48 St. Audreys Close, Hatfield, Hertfordshire AL10 8UW (GB). MATASSA, Victor, Giulio (GB/IT); Istituto Ricerche di Biologia Molecolare, Via Pontina Km. 30,600, I-00040 Pomezia (IT). REEVE, Austin, John (GB/GB); Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). STERNFELD, Francine (GB/GB); Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). STREET, Leslie, Joseph (GB/GB); Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).		(74) Agent: THOMPSON, John; Merck & Co., Inc., European Patent Dept., Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). (81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG). Published <i>With international search report.</i>	
(54) Title: AZETIDINE, PYRROLIDINE AND PIPERIDINE DERIVATIVES AS 5HT₁ RECEPTOR AGONISTS			
(57) Abstract A class of substituted azetidine, pyrrolidine and piperidine derivatives are selective agonists of 5-HT ₁ -like receptors, being potent agonists of the human 5-HT _{1D} receptor subtype whilst possessing at least a 10-fold selective affinity for the 5-HT _{1D} receptor subtype relative to the 5-HT _{1A} subtype; they are therefore useful in the treatment and/or prevention of clinical conditions, in particular migraine and associated disorders, for which a subtype-selective agonist of 5-HT _{1D} receptors is indicated, whilst eliciting fewer side-effects, notably adverse cardiovascular events, than those associated with non-subtype-selective 5-HT _{1D} receptor agonists.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

AZETIDINE, PYRROLIDINE AND PIPERIDINE DERIVATIVES AS 5HT₁ RECEPTOR AGONISTS

The present invention relates to a class of substituted azetidine, pyrrolidine and piperidine derivatives which act on 5-hydroxytryptamine (5-HT) receptors, being selective agonists of so-called "5-HT₁-like" receptors. They are therefore useful in the treatment of clinical conditions for which a selective agonist of these receptors is indicated.

It has been known for some time that 5-HT₁-like receptor agonists which exhibit selective vasoconstrictor activity are of use in the treatment of migraine (see, for example, A. Doenicke *et al.*, *The Lancet*, 1988, Vol. 1, 1309-11; and W. Feniuk and P.P.A. Humphrey, *Drug Development Research*, 1992, 26, 235-240).

The human 5-HT₁-like or 5-HT_{1D} receptor has recently been shown by molecular cloning techniques to exist in two distinct subtypes. These subtypes have been termed 5-HT_{1Dα} (or 5-HT_{1D-1}) and 5-HT_{1Dβ} (or 5-HT_{1D-2}), and their amino acid sequences are disclosed and claimed in WO-A-91/17174.

The 5-HT_{1Dα} receptor subtype in humans is believed to reside on sensory terminals in the dura mater. Stimulation of the 5-HT_{1Dα} subtype inhibits the release of inflammatory neuropeptides which are thought to contribute to the headache pain of migraine. The human 5-HT_{1Dβ} receptor subtype, meanwhile, is located predominantly on the blood vessels and in the brain, and hence may play a part in mediating constriction of cerebral and coronary arteries, as well as CNS effects.

Administration of the prototypical 5-HT_{1D} agonist sumatriptan (GR43175) to humans is known to give rise at therapeutic doses to certain adverse cardiovascular events (see, for example, F. Willett *et al.*, *Br. Med. J.*, 1992, 304, 1415; J.P. Ottervanger *et al.*, *The Lancet*, 1993, 341, 861-2; and D.N. Bateman, *The Lancet*, 1993, 341, 221-4). Since sumatriptan barely discriminates between the human 5-HT_{1Dα} and 5-HT_{1Dβ} receptor subtypes (cf. WO-A-91/17174, Table 1), and since it is the blood vessels

with which the 5-HT_{1D β} subtype is most closely associated, it is believed that the cardiovascular side-effects observed with sumatriptan can be attributed to stimulation of the 5-HT_{1D β} receptor subtype. It is accordingly considered (cf. G.W. Rebeck *et al.*, *Proc. Natl. Acad. Sci. USA*, 1994, 91, 3666-9) that compounds which can interact selectively with the 5-HT_{1D α} receptor subtype, whilst having a less pronounced action at the 5-HT_{1D β} subtype, might be free from, or at any rate less prone to, the undesirable cardiovascular and other side-effects associated with non-subtype-selective 5-HT_{1D} receptor agonists, whilst at the same time maintaining a beneficial level of anti-migraine activity.

The compounds of the present invention, being selective 5-HT₁-like receptor agonists, are accordingly of benefit in the treatment of migraine and associated conditions, e.g. cluster headache, chronic paroxysmal hemicrania, headache associated with vascular disorders, tension headache and paediatric migraine. In particular, the compounds according to this invention are potent agonists of the human 5-HT_{1D α} receptor subtype. Moreover, the compounds in accordance with this invention have been found to possess at least a 10-fold selective affinity for the 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype, and they can therefore be expected to manifest fewer side-effects than those associated with non-subtype-selective 5-HT_{1D} receptor agonists.

Several distinct classes of heteroaromatic compounds based on *inter alia* a substituted tryptamine ring system are described in published International patent applications 91/18897, 94/02460 and 94/02477. The compounds described therein are stated to be agonists of 5-HT₁-like receptors, and accordingly to be of particular use in the treatment of migraine and associated conditions. None of these publications, however, discloses or even suggests the substituted azetidine, pyrrolidine and piperidine derivatives provided by the present invention.


In EP-A-0548813 is described a series of alkoxypyridin-4-yl and alkoxypyrimidin-4-yl derivatives of indol-3-ylalkylpiperazines which are

alleged to provide treatment of vascular or vascular-related headaches, including migraine. There is, however, no disclosure nor any suggestion in EP-A-0548813 of replacing the precisely substituted piperazine moiety described therein with a substituted azetidine, pyrrolidine or piperidine moiety.

Moreover, nowhere in the prior art available to date is there any disclosure of a subtype-selective 5-HT_{1D} receptor agonist having a 5-HT_{1D α} receptor binding affinity (IC₅₀) below 50 nM and at least a 10-fold selective affinity for the 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype.

The compounds according to the present invention are subtype-selective 5-HT_{1D} receptor agonists having a human 5-HT_{1D α} receptor binding affinity (IC₅₀) below 50 nM, typically below 10 nM and preferably below 1 nM; and at least a 10-fold selective affinity, typically at least a 50-fold selective affinity and preferably at least a 100-fold selective affinity, for the human 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype.

The present invention provides a compound of formula I, or a salt or prodrug thereof:



(I)


20

wherein


Z represents hydrogen, halogen, cyano, nitro, trifluoromethyl, -OR⁵, -OCOR⁵, -OCONR⁵R⁶, -OCH₂CN, -OCH₂CONR⁵R⁶, -SR⁵, -SOR⁵, -SO₂R⁵, -SO₂NR⁵R⁶, -NR⁵R⁶, -NR⁵COR⁶, -NR⁵CO₂R⁶, -NR⁵SO₂R⁶, -COR⁵, -CO₂R⁵, -CONR⁵R⁶, or a group of formula (a), (b), (c) or (d):

25


- 4 -




(a)



(b)



(c)



(d)

in which the asterisk * denotes a chiral centre;

X represents oxygen, sulphur, -NH- or methylene;

5 Y represents oxygen or sulphur;

E represents a chemical bond or a straight or branched alkylene chain containing from 1 to 4 carbon atoms;

Q represents a straight or branched alkylene chain containing from 1 to 4 carbon atoms, optionally substituted in any position by a hydroxy

10 group;

U represents nitrogen or C-R²;

V represents oxygen, sulphur or N-R³;

R², R³ and R⁴ independently represent hydrogen or C₁₋₆ alkyl;

R⁵ and R⁶ independently represent hydrogen, C₁₋₆ alkyl,

15 trifluoromethyl, phenyl, methylphenyl, or an optionally substituted aryl(C₁₋₆)alkyl or heteroaryl(C₁₋₆)alkyl group; or R⁵ and R⁶, when linked through a nitrogen atom, together represent the residue of an optionally substituted azetidine, pyrrolidine, piperidine, morpholine or piperazine ring;

20 M represents the residue of an azetidine, pyrrolidine or piperidine ring;

R represents a group of formula -W-R¹;

W represents a chemical bond or a straight or branched alkylene chain containing from 1 to 4 carbon atoms;

25 R¹ represents -OR^x, -SR^x or -NR^xR^y;

R^x and R^y independently represent hydrogen, hydrocarbon or a heterocyclic group, or R^x and R^y together represent a C₂₋₆ alkylene group; and

5 R^z represents hydrogen, hydroxy, hydrocarbon or a heterocyclic group.

The present invention also provides compounds of formula I above, and salts and prodrugs thereof, wherein R⁵ and R⁶ independently represent hydrogen, C₁₋₆ alkyl, trifluoromethyl, methylphenyl, or an optionally substituted aryl(C₁₋₆)alkyl or heteroaryl(C₁₋₆)alkyl group.

10 When R⁵ and R⁶, when linked through a nitrogen atom, together represent the residue of an azetidine, pyrrolidine, piperidine, morpholine or piperazine ring, this ring may be unsubstituted or substituted by one or more substituents. Examples of suitable substituents include C₁₋₆ alkyl, aryl(C₁₋₆)alkyl, C₁₋₆ alkoxy, C₂₋₆ alkoxycarbonyl and C₁₋₆
15 alkylaminocarbonyl. Typical substituents include methyl, benzyl, methoxy, methoxycarbonyl, ethoxycarbonyl and methylaminocarbonyl. In particular, where R⁵ and R⁶ together represent the residue of a piperazine ring, this ring is preferably substituted on the distal nitrogen atom by a C₂₋₆ alkoxycarbonyl moiety such as methoxycarbonyl or ethoxycarbonyl.

20 For use in medicine, the salts of the compounds of formula I will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which
25 may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or
30 phosphoric acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof

may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts.

The term "hydrocarbon" as used herein includes straight-chained,
5 branched and cyclic groups containing up to 18 carbon atoms, suitably up to 15 carbon atoms, and conveniently up to 12 carbon atoms. Suitable hydrocarbon groups include C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl(C₁₋₆)alkyl, aryl and aryl(C₁₋₆)alkyl.

The expression "a heterocyclic group" as used herein includes cyclic
10 groups containing up to 18 carbon atoms and at least one heteroatom preferably selected from oxygen, nitrogen and sulphur. The heterocyclic group suitably contains up to 15 carbon atoms and conveniently up to 12 carbon atoms, and is preferably linked through carbon. Examples of suitable heterocyclic groups include C₃₋₇ heterocycloalkyl, C₃₋₇
15 heterocycloalkyl(C₁₋₆)alkyl, heteroaryl and heteroaryl(C₁₋₆)alkyl groups.

Suitable alkyl groups include straight-chained and branched alkyl groups containing from 1 to 6 carbon atoms. Typical examples include methyl and ethyl groups, and straight-chained or branched propyl and butyl groups. Particular alkyl groups are methyl, ethyl, n-propyl,
20 isopropyl and t-butyl.

Suitable alkenyl groups include straight-chained and branched alkenyl groups containing from 2 to 6 carbon atoms. Typical examples include vinyl and allyl groups.

Suitable alkynyl groups include straight-chained and branched
25 alkynyl groups containing from 2 to 6 carbon atoms. Typical examples include ethynyl and propargyl groups.

Suitable cycloalkyl groups include groups containing from 3 to 7 carbon atoms. Particular cycloalkyl groups are cyclopropyl and cyclohexyl.

30 Particular aryl groups include phenyl and naphthyl.

Particular aryl(C₁₋₆)alkyl groups include benzyl, phenylethyl, phenylpropyl and naphthylmethyl.

Suitable heterocycloalkyl groups include azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny groups.

5 Suitable heteroaryl groups include pyridyl, quinolyl, isoquinolyl, pyridaziny, pyrimidiny, pyraziny, pyranly, furyl, benzofuryl, dibenzofuryl, thienyl, benzthienyl, pyrroly, indolyl, pyrazolyl, indazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl and tetrazolyl groups.

10 The expression "heteroaryl(C₁₋₆)alkyl" as used herein includes furylmethyl, furylethyl, thienylmethyl, thienylethyl, oxazolylmethyl, oxazolylethyl, thiazolylmethyl, thiazolylethyl, imidazolylmethyl, imidazolylethyl, oxadiazolylmethyl, oxadiazolylethyl, thiadiazolylmethyl, thiadiazolylethyl, triazolylmethyl, triazolylethyl, tetrazolylmethyl, 15 tetrazolylethyl, pyridylmethyl, pyridylethyl, pyrimidinylmethyl, pyrazinylmethyl, quinolylmethyl and isoquinolylmethyl.

The hydrocarbon and heterocyclic groups, as well as the aryl(C₁₋₆)alkyl or heteroaryl(C₁₋₆)alkyl groups R⁵ and/or R⁶, may in turn be optionally substituted by one or more groups selected from C₁₋₆ alkyl, 20 adamantyl, phenyl, halogen, C₁₋₆ haloalkyl, C₁₋₆ aminoalkyl, trifluoromethyl, hydroxy, C₁₋₆ alkoxy, aryloxy, keto, C₁₋₃ alkylenedioxy, nitro, cyano, carboxy, C₂₋₆ alkoxy carbonyl, C₂₋₆ alkoxy carbonyl(C₁₋₆)alkyl, C₂₋₆ alkyl carbonyloxy, aryl carbonyloxy, C₂₋₆ alkyl carbonyl, aryl carbonyl, C₁₋₆ alkylthio, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, arylsulphonyl, 25 -NR^vR^w, -NR^vCOR^w, -NR^vCO₂R^w, -NR^vSO₂R^w, -CH₂NR^vSO₂R^w, -NHCONR^vR^w, -CONR^vR^w, -SO₂NR^vR^w and -CH₂SO₂NR^vR^w, in which R^v and R^w independently represent hydrogen, C₁₋₆ alkyl, aryl or aryl(C₁₋₆)alkyl, or R^v and R^w together represent a C₂₋₆ alkylene group.

30 When R^v and R^v, or R^v and R^w, together represent a C₂₋₆ alkylene group, this group may be an ethylene, propylene, butylene,

pentamethylene or hexamethylene group, preferably butylene or pentamethylene.


The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, especially fluorine.

5 The present invention includes within its scope prodrugs of the compounds of formula I above. In general, such prodrugs will be functional derivatives of the compounds of formula I which are readily convertible *in vivo* into the required compound of formula I. Conventional procedures for the selection and preparation of suitable prodrug
10 derivatives are described, for example, in *Design of Prodrugs*, ed. H. Bundgaard, Elsevier, 1985.

Where the compounds according to the invention have at least one asymmetric centre, they may accordingly exist as enantiomers. For example, the compounds of formula I above wherein Z represents a group
15 of formula (b) or (c) have a chiral centre denoted by the asterisk *, which may accordingly be in the (*R*) or (*S*) configuration. Where the compounds according to the invention possess two or more asymmetric centres, they may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed
20 within the scope of the present invention.

In particular, where M represents the residue of a pyrrolidine ring, and the substituent R is attached to the 2-position thereof, then the absolute stereochemical configuration of the carbon atom at the point of attachment of the moiety R is preferably as depicted in structure IA as
25 follows:


- 9 -



(IA)

wherein Z, E, Q, U, V, R and R^{*} are as defined above.

- Moreover, where M represents the residue of a pyrrolidine ring, and
 5 the substituent R is attached to the 3-position thereof, then the absolute stereochemical configuration of the carbon atom at the point of attachment of the moiety R is preferably as depicted in structure IB as follows:



(IB)

wherein Z, E, Q, U, V, R and R^{*} are as defined above.



- Where E, Q and W, which may be the same or different, represent straight or branched alkylene chains, these may be, for example,
 15 methylene, ethylene, 1-methylethylene, propylene, 2-methylpropylene or butylene. In addition, the alkylene chain Q may be substituted in any position by a hydroxy group giving rise, for example, to a 2-hydroxypropylene or 2-hydroxymethyl-propylene chain Q. Moreover, E and W may each independently represent a chemical bond. Where E
 20 represents a chemical bond, the moiety Z is attached directly to the benzo moiety of the central fused bicyclic heteroaromatic ring system. Similarly, where W represents a chemical bond, the substituent R¹ is attached

directly to the azetidine, pyrrolidine or piperidine ring of which M is the residue.

Suitably, E represents a chemical bond or a methylene linkage.


Suitably, Q represents an ethylene or propylene linkage.

- 5 The compound of formula I in accordance with the present invention is suitably an indole, benzofuran or benzthiophene derivative of formula IC, or an indazole derivative of formula ID:



10

wherein Z, E, Q, V, M, R, R¹, R² and R³ are as defined above. Preferably, the compounds according to the invention are indole derivatives of formula IE:



15

wherein Z, E, Q, M, R, R¹, R² and R³ are as defined above, in particular wherein R² and R³ are both hydrogen.

Suitably, W represents a chemical bond or a methylene linkage.

Suitably, R^x and R^y independently represent hydrogen, C₁₋₆ alkyl, aryl, aryl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₆)alkyl, any of which groups may be optionally substituted by one or more substituents selected
5 typically from halogen, hydroxy, C₁₋₆ alkoxy, amino, C₂₋₆ alkylcarbonylamino, C₁₋₆ alkylsulphonylamino and C₁₋₆ alkylaminosulphonylmethyl. Particular values of R^x and R^y include hydrogen, methyl, benzyl, fluorobenzyl, methoxy-benzyl, acetylamino-benzyl, 1-phenylethyl, 2-phenylethyl, 2-hydroxy-1-phenylethyl, 1-
10 (acetylamino-phenyl)ethyl, 2-(acetylamino-phenyl)ethyl, 1-hydroxy-3-phenylprop-2-yl, 1-hydroxy-1-phenylprop-2-yl, furylmethyl, thienylmethyl and pyridylmethyl.

Suitable values for the substituent R¹ include hydroxy, benzyloxy, methoxy-benzyloxy, pyridylmethoxy, amino, methylamino, benzylamino,
15 N-(acetylamino-benzyl)-amino, N-(1-phenylethyl)-amino, N-(2-phenylethyl)-amino, N-(2-hydroxy-1-phenylethyl)-amino, N-[1-(acetylamino-phenyl)ethyl]-amino, N-[2-(acetylamino-phenyl)ethyl]-amino, N-(1-hydroxy-3-phenylprop-2-yl)-amino, N-(1-hydroxy-1-phenylprop-2-yl)-amino, N-(furylmethyl)-amino, N-(pyridylmethyl)-amino,
20 dimethylamino, N-benzyl-N-methylamino, N-fluorobenzyl-N-methylamino, N-(acetylamino-benzyl)-N-methylamino, N-methyl-N-(1-phenylethyl)-amino, N-(2-hydroxy-1-phenylethyl)-N-methylamino, N-[2-(acetylamino-phenyl)ethyl]-N-methylamino and N-methyl-N-(thienylmethyl)-amino.

Particular values of the group R include hydroxy, benzyloxy, benzyloxymethyl, methoxy-benzyloxy, pyridylmethoxy, benzylamino, benzylaminomethyl, N-(acetylamino-benzyl)-amino, N-(acetylamino-benzyl)-aminomethyl, N-(1-phenylethyl)-amino, N-(1-phenylethyl)-aminomethyl, N-(2-hydroxy-1-phenylethyl)-amino, N-(2-hydroxy-1-phenylethyl)-aminomethyl, N-[1-(acetylamino-phenyl)ethyl]-amino, N-[1-(acetylamino-phenyl)ethyl]-aminomethyl, N-[2-(acetylamino-
30 (acetylamino-phenyl)ethyl]-aminomethyl, N-[2-(acetylamino-

phenyl)ethyl]-amino, N-(1-hydroxy-3-phenylprop-2-yl)-amino, N-(1-hydroxy-1-phenylprop-2-yl)-amino, N-(furylmethyl)-aminomethyl, N-(pyridylmethyl)-aminomethyl, N-benzyl-N-methylamino, N-benzyl-N-methyl-aminomethyl, N-fluorobenzyl-N-methyl-aminomethyl, N-
5 (acetyl-amino-benzyl)-N-methyl-aminomethyl, N-methyl-N-(1-phenylethyl)-aminomethyl, N-(2-hydroxy-1-phenylethyl)-N-methylamino, N-[2-(acetyl-amino-phenyl)ethyl]-N-methylamino and N-methyl-N-(thienylmethyl)amino.

Suitable values of R^a include hydrogen, hydroxy and benzyl,
10 especially hydrogen.

Suitably, R² and R³ independently represent hydrogen or methyl, especially hydrogen.

Suitably, R⁴ represents hydrogen or methyl.


Suitably, R⁵ and R⁶ are independently selected from hydrogen,
15 methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, trifluoromethyl, phenyl, methylphenyl (especially 4-methylphenyl), benzyl and phenethyl.

Suitably, the substituent Z represents hydrogen, fluoro, cyano, hydroxy, methoxy, ethoxy, benzyloxy, methylamino-carbonyloxy, cyano-methoxy, aminocarbonyl-methoxy, methylsulphonyl, phenylsulphonyl,
20 aminosulphonyl, N-methylamino-sulphonyl, N,N-dimethylamino-sulphonyl, amino, formylamino, acetyl-amino, trifluoromethyl-carbonylamino, benzyloxy-carbonylamino, methyl-sulphonylamino, ethyl-sulphonylamino, methylphenyl-sulphonylamino, N-methyl-(N-methylsulphonyl)-amino, N-methyl-(N-ethylsulphonyl)-amino, N-methyl-
25 (N-trifluoromethylsulphonyl)-amino, N-ethyl-(N-methylsulphonyl)-amino, N-benzyl-(N-methylsulphonyl)-amino, N-benzyl-(N-ethylsulphonyl)-amino, acetyl, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl, methylaminocarbonyl, ethylaminocarbonyl, propylaminocarbonyl, butylaminocarbonyl, benzylaminocarbonyl or phenethyl-aminocarbonyl; or
30 a group of formula (a), (b), (c) or (d) as defined above.

In a particular embodiment, Z represents $-\text{SO}_2\text{NR}^5\text{R}^6$ in which R^5 and R^6 are as defined above. In a subset of this embodiment, R^5 and R^6 independently represent hydrogen or C_{1-6} alkyl, especially hydrogen or methyl. Particular values of Z in this context include aminosulphonyl, N-methylamino-sulphonyl and N,N-dimethylamino-sulphonyl, especially N-methylamino-sulphonyl.

In another embodiment, Z represents a group of formula (b) in which R^4 is hydrogen or methyl. In a subset of this embodiment, X and Y both represent oxygen. In a particular aspect of this subset, the chiral centre denoted by the asterisk * is in the (S) configuration.

A particular sub-class of compounds according to the invention is represented by the compounds of formula IIA, and salts and prodrugs thereof:



wherein

m is zero, 1, 2 or 3, preferably zero or 1;

n is 2, 3 or 4, preferably 2 or 3;

p is zero, 1 or 2;

R^5 and R^6 are as defined with reference to formula I above;

W^1 represents oxygen, sulphur or N-R^{12} ; and

R^{11} and R^{12} independently represent hydrogen, C_{1-6} alkyl, aryl, aryl(C_{1-6})alkyl, heteroaryl or heteroaryl(C_{1-6})alkyl, any of which groups may be optionally substituted.

Examples of suitable optional substituents on the groups R^{11} and R^{12} include halogen, cyano, trifluoromethyl, hydroxy, C_{1-6} alkoxy, C_{2-6} alkylcarbonyl, amino, C_{1-6} alkylamino, di(C_{1-6})alkylamino, C_{2-6} alkylcarbonylamino, C_{1-6} alkylsulphonylamino and C_{1-6}

5 alkylaminosulphonylmethyl.


Particular values of R^5 and R^6 with reference to formula IIA above include hydrogen and C_{1-6} alkyl, especially hydrogen or methyl. Suitably, one of R^5 and R^6 represents hydrogen and the other represents hydrogen or methyl.

10 Particular values of R^{11} and R^{12} include hydrogen, methyl, benzyl, fluorobenzyl, methoxy-benzyl, acetylamino-benzyl, 1-phenylethyl, 2-phenylethyl, 2-hydroxy-1-phenylethyl, 1-(acetylamino-phenyl)ethyl, 2-(acetylamino-phenyl)ethyl, 1-hydroxy-3-phenylprop-2-yl, 1-hydroxy-1-phenylprop-2-yl, furylmethyl, thienylmethyl and pyridylmethyl.

15 Typically, R^{11} represents benzyl, fluorobenzyl, 1-phenylethyl or 2-hydroxy-1-phenylethyl.

Typically, R^{12} is hydrogen or methyl.

Another sub-class of compounds according to the invention is represented by the compounds of formula IIB, and salts and prodrugs thereof:




wherein the asterisk * denotes a chiral centre;

25 R^4 and Y are as defined with reference to formula I above; and

m, n, p, W¹ and R¹¹ are as defined with reference to formula IIA above.

A further sub-class of compounds according to the invention is represented by the compounds of formula IIC, and salts and prodrugs thereof:



wherein the asterisk * denotes a chiral centre;

R⁴ and Y are as defined with reference to formula I above; and
m, n, p, W¹ and R¹¹ are as defined with reference to formula IIA above.

In relation to formula IIB and IIC above, the chiral centre denoted by the asterisk * is suitably in the (S) configuration.

Specific compounds within the scope of the present invention include:

- (3S)-3-(N-benzyl)aminomethyl-1-[2-(5-(N-methyl)-aminosulphonylmethyl)-1H-indol-3-yl]ethyl]pyrrolidine;
- (3S)-3-(N-benzyl)aminomethyl-1-[2-(5-(aminosulphonylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine;
- (3S)-3-(N-benzyl)aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine;
- (3S)-3-[N-(R)-α-(hydroxymethyl)benzyl]aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine;
- (3S)-3-[N-(S)-α-methylbenzyl]aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine;

- 4-[*N*-(*R*)- α -(hydroxymethyl)benzyl]amino-(*S*)-1-[3-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)propyl]piperidine;
(3*S*)-3-(*N*-benzyl-*N*-methyl)aminomethyl-(*S*)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
5 (3*R*)-3-[*N*-(*S*)- α -methylbenzyl-*N*-methyl]aminomethyl-(*S*)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
(3*R*)-3-[*N*-(*S*)- α -methylbenzyl-*N*-methyl]aminomethyl-(*S*)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
(3*S*)-3-[*N*-(4-fluorobenzyl)-*N*-methyl]aminomethyl-(*S*)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
10 and salts and prodrugs thereof.


The invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier. Preferably these compositions are in
15 unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. For preparing solid
20 compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing
25 a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage
30 forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type

described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of the active ingredient. The tablets or pills of the novel composition can be coated or
5 otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and
10 permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.


15 The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar
20 pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

In the treatment of migraine, a suitable dosage level is about 0.01
25 to 250 mg/kg per day, preferably about 0.05 to 100 mg/kg per day, and especially about 0.05 to 5 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day.

The compounds according to the invention wherein U represents C-R² and V represents N-R³, corresponding to the indole derivatives of
30 formula IE as defined above, may be prepared by a process which comprises reacting a compound of formula III:



wherein Z and E are as defined above; with a compound of formula IV, or
 5 a carbonyl-protected form thereof:




wherein R², Q, M, R and R' are as defined above; followed, where
 10 required, by N-alkylation by standard methods to introduce the moiety R³.

The reaction between compounds III and IV, which is an example of the well-known Fischer indole synthesis, is suitably carried out by heating the reagents together under mildly acidic conditions, e.g. 4% sulphuric acid at reflux.

15 Suitable carbonyl-protected forms of the compounds of formula IV include the dimethyl acetal or ketal derivatives.


The Fischer reaction between compounds III and IV may be carried out in a single step, or may proceed via an initial non-cyclising step at a lower temperature to give an intermediate of formula V:




(V)

wherein Z, E, Q, R², M, R and R^{*} are as defined above; followed by cyclisation using a suitable reagent, e.g. a polyphosphate ester.

- 5 The intermediates of formula IV, or carbonyl-protected forms thereof, may be prepared by reacting a compound of formula VI, or a carbonyl-protected form thereof, with a compound of formula VII:



(VI)



(VII)

10


wherein Q, R², M, R and R^{*} are as defined above, and L¹ represents a suitable leaving group.

The leaving group L¹ is suitably a halogen atom, e.g. chlorine or bromine.

- 15 Where L¹ represents a halogen atom, the reaction between compounds VI and VII is conveniently effected by stirring the reactants under basic conditions in a suitable solvent, for example sodium carbonate or potassium carbonate in 1,2-dimethoxyethane or *N,N*-dimethylformamide, or triethylamine in tetrahydrofuran or acetonitrile,
- 20 optionally in the presence of catalytic sodium iodide.

In an alternative procedure, the compounds according to the invention may be prepared by a process which comprises reacting a

compound of formula VII as defined above with a compound of formula VIII:



(VIII)


5

wherein Z, E, Q, U and V are as defined above, and L² represents a suitable leaving group.

The leaving group L² is suitably an alkylsulphonyloxy or arylsulphonyloxy group, e.g. methanesulphonyloxy (mesyloxy) or *p*-toluenesulphonyloxy (tosyloxy).

Where L² represents an alkylsulphonyloxy or arylsulphonyloxy group, the reaction between compounds VII and VIII is conveniently carried out in a suitable solvent such as isopropanol/acetonitrile *N,N*-dimethylformamide or 1,2-dimethoxyethane, typically in the presence of a base such as sodium carbonate or potassium carbonate, optionally in the presence of catalytic sodium iodide.

In one representative approach, the compounds of formula VIII wherein U represents CH, V represents NH and L² represents a mesyloxy or tosyloxy group may be prepared by the sequence of steps illustrated in the following reaction scheme (cf. Larock and Yum, *J. Am. Chem. Soc.*, 1991, 113, 6689):



wherein Z, E and Q are as defined above, L^3 represents mesyloxy or tosyloxy, and TES is an abbreviation for triethylsilyl.

- 5 In Step 1 of the reaction scheme, the aniline derivative IX is treated with iodine monochloride, typically in acetonitrile, in order to introduce an iodine atom *ortho* to the amine moiety. Step 2 involves a palladium-mediated coupling reaction with the protected acetylene derivative $\text{TES-C}\equiv\text{C-Q-OTES}$, typically using palladium acetate and
- 10 triphenylphosphine in the presence of lithium chloride and sodium carbonate, suitably in *N,N*-dimethylformamide at an elevated temperature. This is followed in Step 3 by removal of the TES moiety, ideally in refluxing methanolic hydrochloric acid; followed in turn by mesylation or tosylation, suitably by using mesyl chloride or tosyl chloride
- 15 respectively in the presence of a base such as triethylamine or pyridine, typically in dichloromethane/acetonitrile.


In another representative approach, the compounds of formula VIII wherein U represents CH, V represents NH, Q represents a propylene chain and L^2 represents a mesyloxy or tosyloxy group may be prepared by

20 reacting 3,4-dihydro-2*H*-pyran with a compound of formula III as defined above or a salt thereof, under a variant of the Fischer reaction conditions as described above for the reaction between compounds III and IV; followed by mesylation or tosylation of the 3-hydroxypropyl-indole

derivative thereby obtained, typically by treatment with mesyl chloride or tosyl chloride under standard conditions.

The Fischer reaction with 3,4-dihydro-2*H*-pyran is suitably brought about by heating an acid addition salt of the hydrazine derivative III,
 5 typically the hydrochloride salt, in an inert solvent such as dioxan, at the reflux temperature of the solvent.

In a further procedure, the compounds according to the invention wherein U represents nitrogen and V represents N-R³, corresponding to the indazole derivatives of formula ID as defined above, may be prepared
 10 by a process which comprises cyclising a compound of formula X:




(X)

wherein Z, E, Q, M, R and R^{*} are as defined above, and D¹ represents a
 15 readily displaceable group; followed, where required, by N-alkylation by standard methods to introduce the moiety R³.

The cyclisation of compound X is conveniently achieved in a suitable organic solvent at an elevated temperature, for example in a mixture of *m*-xylene and 2,6-lutidine at a temperature in the region of
 20 140°C.


The readily displaceable group D¹ in the compounds of formula X suitably represents a C₁₋₄ alkanoyloxy group, preferably acetoxy. Where D¹ represents acetoxy, the desired compound of formula X may be conveniently prepared by treating a carbonyl compound of formula XI:



(XI)

wherein Z, E, Q, M, R and R^{*} are as defined above; or a protected derivative thereof, preferably the N-formyl protected derivative; with
 5 hydroxylamine hydrochloride, advantageously in pyridine at the reflux temperature of the solvent; followed by acetylation with acetic anhydride, advantageously in the presence of a catalytic quantity of 4-
 dimethylaminopyridine, in dichloromethane at room temperature.

The N-formyl protected derivatives of the intermediates of formula
 10 XI may conveniently be prepared by ozonolysis of the corresponding indole derivative of formula XII:




(XII)

15 wherein Z, E, Q, M, R and R^{*} are as defined above; followed by a reductive work-up, advantageously using dimethylsulphide.

The indole derivatives of formula XII may be prepared by methods analogous to those described in the accompanying Examples, or by procedures well known from the art.

20 In a still further procedure, the compounds according to the invention wherein U represents C-R² and V represents oxygen or sulphur,

corresponding to the benzofuran or benzthiophene derivatives of formula IC wherein V is oxygen or sulphur respectively, may be prepared by a process which comprises cyclising a compound of formula XIII:




(XIII)


wherein Z, E, Q, R², M, R and R* are as defined above, and V¹ represents oxygen or sulphur.

The cyclisation of compound XIII is conveniently effected by using polyphosphoric acid or a polyphosphate ester, advantageously at an elevated temperature.

The compounds of formula XIII may be prepared by reacting a compound of formula XIV with a compound of formula XV:



(XIV)



(XV)

wherein Z, E, Q, R², V¹, M, R and R* are as defined above, and Hal represents a halogen atom.

The reaction is conveniently effected in the presence of a base such as sodium hydroxide.

The hydroxy and mercapto derivatives of formula XIV may be prepared by a variety of methods which will be readily apparent to those skilled in the art.

The hydrazine derivatives of formula III above may be prepared by methods analogous to those described in EP-A-0548813 and WO-A-91/18897, as also may the aniline derivatives of formula IX.

Where they are not commercially available, the starting materials of formula VI, VII and XV may be prepared by the methods described in the accompanying Examples, or by analogous procedures which will be apparent to those skilled in the art.

It will be understood that any compound of formula I initially obtained from any of the above processes may, where appropriate, subsequently be elaborated into a further compound of formula I by techniques known from the art. For example, a compound of formula I wherein R^x is benzyl initially obtained may be converted into a compound of formula I wherein R^x is hydrogen typically by conventional catalytic hydrogenation, or by transfer hydrogenation using a hydrogenation catalyst such as palladium on charcoal in the presence of a hydrogen donor such as ammonium formate. Moreover, a compound of formula I wherein R^1 is hydroxy initially obtained may be converted into the corresponding carbonyl compound (aldehyde or ketone) by treatment with a conventional oxidising agent such as sulphur trioxide-pyridine complex; the resulting carbonyl compound may then be converted in turn into a compound of formula I wherein R^1 represents $-NHR^y$, suitably by a standard reductive amination procedure which comprises treating the carbonyl compound with the appropriate amine of formula R^y-NH_2 in the presence of a suitable reducing agent, typically sodium cyanoborohydride. Similarly, a compound of formula I wherein R^1 represents $-NHR^y$ initially obtained may be converted into a further compound of formula I wherein R^1 represents $-NR^xR^y$, in which R^x corresponds to the group $-CH_2R^z$, suitably by a reductive amination procedure which comprises treating the compound of formula I wherein R^1 represents $-NHR^y$ with the appropriate aldehyde of formula R^z-CHO in the presence of a reducing agent such as sodium cyanoborohydride. In addition, a compound of formula I wherein

R³ is hydrogen initially obtained may be converted into a compound of formula I wherein R³ represents C₁₋₆ alkyl by standard alkylation techniques, for example by treatment with an alkyl iodide, e.g. methyl iodide, typically under basic conditions, e.g. sodium hydride in
5 dimethylformamide, or triethylamine in acetonitrile.

Where the above-described processes for the preparation of the compounds according to the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The novel compounds may be
10 prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The novel compounds may, for example, be resolved into their component enantiomers by standard techniques such as preparative HPLC, or the formation of diastereomeric pairs by salt formation with an optically active acid, such
15 as (-)-di-*p*-toluoyl-d-tartaric acid and/or (+)-di-*p*-toluoyl-l-tartaric acid, followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

20 During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene &
25 P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The following Examples illustrate the preparation of compounds according to the invention.

30 The compounds in accordance with the present invention potently and selectively bind to the 5-HT_{1Dα} receptor subtype, inhibit forskolin-

stimulated adenylyl cyclase activity, and stimulate [³⁵S]-GTP γ S binding to membranes from clonal cell lines expressing human cloned receptors.

5-HT_{1D α} /5-HT_{1D β} Radioligand Binding

5

Chinese hamster ovary (CHO) clonal cell lines expressing the human 5-HT_{1D α} and 5-HT_{1D β} receptors were harvested in PBS and homogenised in ice cold 50 mM Tris-HCl (pH 7.7 at room temperature) with a Kinematica polytron and centrifuged at 48,000g at 4°C for 11 min.

10 The pellet was then resuspended in 50 mM Tris-HCl followed by a 10 min incubation at 37°C. Finally the tissue was recentrifuged at 48,000g, 4°C for 11 min and the pellet resuspended, in assay buffer (composition in mM: Tris-HCl 50, pargyline 0.01, CaCl₂ 4; ascorbate 0.1%; pH 7.7 at room temperature) to give the required volume immediately prior to use (0.2 mg

15 protein/ml). Incubations were carried out for 30 min at 37°C in the presence of 0.02-150 nM [³H]-5-HT for saturation studies or 2-5 nM [³H]-5-HT for displacement studies. The final assay volume was 1 ml. 5-HT (10 μ M) was used to define non-specific binding. The reaction was initiated by the addition of membrane and was terminated by rapid filtration

20 through Whatman GF/B filters (presoaked in 0.3% PEI/ 0.5% Triton X) followed by 2 x 4 ml washings with 50 mM Tris-HCl. The radioactive filters were then counted on a LKB beta or a Wallac beta plate counter. Binding parameters were determined by non-linear, least squares regression analysis using an iterative curve fitting routine, from which

25 IC₅₀ (the molar concentration of compound necessary to inhibit binding by 50%) values could be calculated for each test compound. The IC₅₀ values for binding to the 5-HT_{1D α} receptor subtype obtained for the compounds of the accompanying Examples were below 50 nM in each case.

30 Furthermore, the compounds of the accompanying Examples were all found to possess a selective affinity for the 5-HT_{1D α} receptor subtype of at least 10-fold relative to the 5-HT_{1D β} subtype.

5-HT_{1D α} /5-HT_{1D β} Adenylyl Cyclase Assay

Studies were performed essentially as described in *J. Pharmacol. Exp. Ther.*, 1986, 238, 248. CHO clonal cell lines expressing the human
5 cloned 5-HT_{1D α} and 5-HT_{1D β} receptors were harvested in PBS and
homogenised, using a motor driven teflon/glass homogeniser, in ice cold
Tris HCl-EGTA buffer (composition in mM: Tris HCl 10, EGTA 1, pH 8.0
at room temperature) and incubated on ice for 30-60 min. The tissue was
then centrifuged at 20,000g for 20 min at 4°C, the supernatant discarded
10 and the pellet resuspended in Tris HCl-EDTA buffer (composition in mM:
Tris HCl 50, EDTA 5, pH 7.6 at room temperature) just prior to assay.
The adenylyl cyclase activity was determined by measuring the conversion
of α -[³³P]-ATP to [³³P]-cyclic AMP. A 10 μ l aliquot of the membrane
suspension was incubated, for 10-15 min, in a final volume of 50 μ l, at
15 30°C, with or without forskolin (10 μ M), in the presence or absence of test
compound. The incubation buffer consisted of 50 mM Tris HCl (pH 7.6 at
room temperature), 100 mM NaCl, 30 μ M GTP, 50 μ M cyclic AMP, 1 mM
dithiothreitol, 1 mM ATP, 5 mM MgCl₂, 1 mM EGTA, 1 mM 3-isobutyl-1-
methylxanthine, 3.5 mM creatinine phosphate, 0.2 mg/ml creatine
20 phosphokinase, 0.5-1 μ Ci α -[³³P]-ATP and 1 nCi [³H]-cyclic AMP. The
incubation was initiated by the addition of membrane, following a 5 min
preincubation at 30°C, and was terminated by the addition of 100 μ l SDS
(composition in mM: sodium lauryl sulphate 2%, ATP 45, cyclic AMP 1.3,
pH 7.5 at room temperature). The ATP and cyclic AMP were separated on
25 a double column chromatography system (*Anal. Biochem.*, 1974, 58, 541).
Functional parameters were determined using a least squares curve
fitting programme ALLFIT (*Am. J. Physiol.*, 1978, 235, E97) from which
E_{max} (maximal effect) and EC₅₀ (the molar concentration of compound
necessary to inhibit the maximal effect by 50%) values were obtained for
30 each test compound. Of those compounds which were tested in this assay,
the EC₅₀ values for the 5-HT_{1D α} receptor obtained for the compounds of the

accompanying Examples were below 500 nM in each case. Moreover, the compounds of the accompanying Examples which were tested were all found to possess at least a 10-fold selectivity for the 5-HT_{1D α} receptor subtype relative to the 5-HT_{1D β} subtype.

5

5-HT_{1D α} /5-HT_{1D β} GTP γ S Binding

Studies were performed essentially as described in *Br. J. Pharmacol.*, 1993, 109, 1120. CHO clonal cell lines expressing the human cloned 5-HT_{1D α} and 5-HT_{1D β} receptors were harvested in PBS and homogenised using a Kinematica polytron in ice cold 20 mM HEPES containing 10 mM EDTA, pH 7.4 at room temperature. The membranes were then centrifuged at 40,000g, 4°C for 15 min. The pellet was then resuspended in ice cold 20 mM HEPES containing 0.1 mM EDTA, pH 7.4 at room temperature and recentrifuged at 40,000g, 4°C for 15-25 minutes. The membranes were then resuspended in assay buffer (composition in mM: HEPES 20, NaCl 100, MgCl₂ 10, pargyline 0.01; ascorbate 0.1%; pH 7.4 at room temperature) at a concentration of 40 μ g protein/ml for the 5-HT_{1D α} receptor transfected cells and 40-50 μ g protein/ml for the 5-HT_{1D β} receptor transfected cells. The membrane suspension was then incubated, in a volume of 1 ml, with GDP (100 μ M for 5-HT_{1D α} receptor transfected cells, 30 μ M for the 5-HT_{1D β} receptor transfected cells) and test compound at 30°C for 20 min and then transferred to ice for a further 15 min. [³⁵S]-GTP γ S was then added at a final concentration of 100 pM and the samples incubated for 30 min at 30°C. The reaction was initiated by the addition of membrane and was terminated by rapid filtration through Whatman GF/B filters and washed with 5 ml water. The radioactive filters were then counted on a LKB beta counter. Functional parameters were determined by a non-linear, least squares regression analysis using an iterative curve fitting routine, from which E_{max} (maximal effect) and EC₅₀ (the molar concentration of compound necessary to inhibit the

maximal effect by 50%) values were obtained for each test compound. Of those compounds which were tested in this assay, the EC₅₀ values for the 5-HT_{1Dα} receptor obtained for the compounds of the accompanying Examples were below 500 nM in each case. Moreover, the compounds of the accompanying Examples which were tested were all found to possess at least a 10-fold selectivity for the 5-HT_{1Dα} receptor subtype relative to the 5-HT_{1Dβ} subtype.

EXAMPLE 1

10

(3S)-3-(N-Benzyl)aminomethyl-1-[2-(5-(N-(methyl)amino-sulphonylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine 2.1 Hydrogen Oxalate 0.7 Diethyl etherate

15 1. Intermediate 1: (3S)-N(H)-3-(N-Benzyl)aminomethylpyrrolidinea) (3R)-N-tert-Butyloxycarbonyl-3-hydroxymethylpyrrolidine

A mixture of (3R)-N-[(R)-1-phenylethyl]-3-

(hydroxymethyl)pyrrolidine (*J. Med. Chem.*, 1990, 33(1), 71; 17.0g, 82.8mmol), di-tert-butylidicarbonate (21.7g, 99.4mmol), Pearlman's catalyst (4.28g, 25% w/w), methanol (300ml) and water (40ml) was hydrogenated on a Parr shake apparatus, at 40 psi, for 2.25h. The mixture was filtered through a pad of celite and the pad washed with ethanol. The combined filtrate and washings were evaporated and the residue chromatographed on silica gel eluting with CH₂Cl₂/MeOH (95:5) to give the title-pyrrolidine (16.73g, 100%), δ (250MHz, D₆-DMSO) 1.39 (9H, s, OC(Me)₃), 1.31-1.64 (2H, m, CH₂), 1.79-1.88 (1H, m, CH), 2.19-2.31 (1H, m, CH of CH₂), 2.95 (1H, dd, J=10.7 and 7.0Hz, CH of CH₂), 3.11-3.35 (4H, m, 2 of CH₂), 4.67 (1H, t, J=5.3Hz, OH).

30

b) (3R)-N-tert-Butyloxycarbonyl-3-methylsulphonyloxy-methylpyrrolidine

A solution of methane sulphonyl chloride (3.37g, 29.39mmol) in CH₂Cl₂ (30ml) was added dropwise to a solution of (3R)-N-tert-butyl-
5 butyloxycarbonyl-3-hydroxymethylpyrrolidine (5.4g, 26.7mmol) and anhydrous triethylamine (2.97g, 29.39mmol), in CH₂Cl₂ (100ml), at -15°C. The mixture was warmed to room temperature and stirred for 16h before adding saturated K₂CO₃ solution (50ml) and diluting with CH₂Cl₂ (100ml). The aqueous was separated and extracted further with CH₂Cl₂ (2x100ml).
10 The combined extracts were dried (Na₂SO₄) and evaporated to give the title-mesylate (7.5g, 100%), δ (250MHz, CDCl₃) 1.46 (9H, s, OC(Me)₃), 1.62-1.84 (1H, m, CH of CH₂), 2.00-2.14 (1H, m, CH of CH₂), 2.58-2.72 (1H, m, CH), 3.04 (3H, s, Me), 3.08-3.62 (4H, m, 2 of CH₂), 4.11-4.33 (2H, m, CH₂OMs).

15

c) (3S)-N-tert-Butyloxycarbonyl-3-N-(benzyl)aminomethylpyrrolidine

A solution of the preceding mesylate (5.0g, 17.90mmol) and benzylamine (9.8ml, 89.7mmol) in toluene (25ml) was heated at reflux for
20 18h. The mixture was evaporated under high vacuum and the residue taken up in ethyl acetate (200ml) and washed with water (x3). The organic layer was dried (MgSO₄) and evaporated and the crude product chromatographed on silica gel eluting with CH₂Cl₂/MeOH (98:2) to give the desired product (4.9g, 94%), δ (250MHz, CDCl₃) 1.45 (9H, s, OC(Me)₃),
25 1.52-1.64 (1H, m, CH), 1.92-2.08 (1H, m, CH of CH₂), 2.27-2.40 (1H, m, CH of CH₂), 2.60-2.68 (2H, m, CH₂), 2.93-3.08 (1H, m, CH of CH₂), 3.18-3.60 (3H, m, CH₂ and CH of CH₂), 3.80 (2H, s, NHCH₂Bn), 7.26-7.36 (5H, m, Ar-H).

30

d) (3S)-N(H)-3-(N-Benzyl)aminomethyl pyrrolidine

A solution of the preceding benzylamine (4.9g, 16.8mmol) in 90% formic acid (90ml) was stirred at room temperature for 18.5h. The reaction was quenched by addition of MeOH and the solvents were removed under vacuum. The residue was dissolved in a minimum volume of H₂O, basified with saturated K₂CO₃ solution and extracted with *n*-butanol (3x100ml). The combined extracts were evaporated *in vacuo* and the inorganics removed by trituration with CH₂Cl₂ and filtration. The filtrate was dried (MgSO₄) and evaporated to give the title-pyrrolidine (3.24g, 100%), δ (360MHz, CDCl₃) 1.42-1.60 (1H, m, CH), 1.94-2.03 (1H, m, CH of CH₂), 2.24-2.36 (1H, m, CH of CH₂), 2.58-2.73 (3H, m, CH₂ and CH of CH₂), 2.94-3.19 (3H, m, CH₂ and CH of CH₂), 3.79 (2H, m, NHCH₂Bn), 7.23-7.35 (5H, m, Ar-H).

2. Intermediate 2: 2-[5-(N-(Methyl)aminosulphonyl-methyl)-1H-indol-3-yl]ethyl alcoholA. 2-Iodo-4-(N-(methyl)aminosulphonylmethyl)phenyl anilinea) 1-(N-(Methyl)aminosulphonylmethyl)-4-nitrobenzene

A mixture of 4-nitrobenzyl bromide (100.0g, 0.46mol), sodium sulphite (84.8g, 0.67mol) and water (316ml) was heated at 90°C for 5h. The solution was cooled and the resultant solid filtered and washed with diethyl ether. The product was dried under vacuum at 60°C (95g, 86%). Phosphorus pentachloride (78g, 0.375mol) was added to sodium 4-nitrobenzylsulphonate (60g, 0.25mol) and the mixture heated at 90°C for 2h. The mixture was cooled and volatile material removed under vacuum. The residue was dissolved in dichloromethane (500ml) and water (150ml). The organic layer was separated, dried over anhydrous sodium sulphate, filtered and evaporated to give 4-nitrobenzyl sulphonyl chloride (48.9g, 83%) which was pure by ¹H NMR. Methylamine gas was bubbled through a solution of 4-nitrobenzyl sulphonyl chloride (37.9g, 0.16mol), in dichloromethane (325ml), until uptake had ceased (0.5h). The resulting

solid was filtered, washed with H₂O and dried under vacuum to give the title-sulphonamide (32.5g, 88%), δ (250MHz, D₆-DMSO) 2.61 (3H, s, Me), 4.55 (2H, s, CH₂), 7.06 (1H, s, NH), 7.66 (2H, d, J=8.7Hz, Ar-H), 8.25 (2H, d, J=8.7Hz, Ar-H).

5

b) 2-Iodo-4-(N-(methyl)aminosulphonylmethyl) phenylaniline

A mixture of the preceding 4-nitro-N-methylbenzenemethane sulphonamide (28.86g, 0.126mol), H₂O (100ml), ethanol (250ml), 5N HCl (25ml) and 10% Pd-C (3.0g) was hydrogenated on a Parr shake apparatus at 50psi for 4h. The catalyst was removed by filtration through celite and the solvents removed under vacuum. The residue was dissolved in water (200ml) and basified with K₂CO₃. The precipitated product was filtered off, washed with water and hexane and dried under vacuum at 45°C to give the desired aniline (21.45g, 85%) which was pure by ¹H NMR. To a stirred suspension of the preceding aniline (21.45g, 0.107mol) in acetonitrile (250ml) was added a solution of iodine monochloride (17.41g, 0.107mol), in acetonitrile (50ml), dropwise over 1h. The mixture was stirred at room temperature for 16h and then partitioned between ethyl acetate (500ml) and 20% aqueous sodium thiosulphate (300ml). The organic layer was separated, washed with H₂O (500ml) and brine (100ml) and dried (Na₂SO₄). The solvent was removed under vacuum and the crude product chromatographed on silica gel eluting with ethyl acetate/hexane (1:1) to give the title-iodoaniline (9.0g, 26%), δ (250MHz, D₆-DMSO) 2.51 (3H, d, J=7.4Hz, Me), 4.11 (2H, s, CH₂), 5.29 (2H, s, NH₂), 6.72 (1H, d, J=8.3Hz, Ar-H), 6.81 (1H, q, J=7.4Hz, NH), 7.06 (1H, dd, J=2.0 and 8.3Hz, Ar-H), 7.53 (1H, d, J=2.0Hz, Ar-H).

B. 1,4-bis-Triethylsilyl-3-butyne-1-ol

n-Butyl lithium (776ml of a 2.5M solution in hexane, 1.94mol) was added over a 2h period to a stirred solution of 3-butyne-1-ol (68g, 0.97mol), in anhydrous THF (1.16l), at -30°C. The mixture was stirred at -30°C for

1h (dianion precipitates out) and triethylsilyl chloride (300g, 1.99mol) was then added dropwise, ensuring that the temperature remained below -20°C. The solution was stirred at -10°C for 1h and then at room temperature for 1h. Hexane (1.36l) and Na₂CO₃ solution (7.0g in 700ml of H₂O) were added to the reaction mixture, at -10°C, and the layers separated. The aqueous layer was extracted with hexane and the combined extracts were washed with water, dried (Na₂SO₄) and evaporated to give the title-bis triethylsilyl butynol (289g, 100%), δ (250MHz, CDCl₃) 0.80-0.94 (12H, m, 6 of SiCH₂CH₃), 1.22-1.32 (18H, m, 6 of SiCH₂CH₃), 2.76 (2H, t, J=7.3Hz, CH₂), 4.00 (2H, t, J=7.3Hz, CH₂).

C. 2-[5-(N-(Methyl)aminosulphonylmethyl)-1H-indol-3-yl]ethyl alcohol

A mixture of 2-iodo-4-(N-(methyl)aminosulphonylmethyl)phenyl aniline (10g, 30.7mmol), 1,4-bis-triethylsilyl-3-butyn-1-ol (10.97g, 36.8mmol) sodium carbonate (16.26g, 153.4mmol) and anhydrous DMF (500ml) was degassed with N₂ for 0.5h. Pd(OAc)₂ (0.7g, 3.1mmol) was added and the mixture heated at 100°C for 6h. The DMF was removed under vacuum and the residue partitioned between EtOAc (250ml) and water (250ml). The solutions were passed through celite to remove insolubles and the aqueous layer separated and extracted further with ethyl acetate (4x200ml). The combined organics were dried (Na₂SO₄) and evaporated. The residue was dissolved in methanol (100ml) and 5N hydrochloric acid (30ml) was added. The mixture was stirred at room temperature for 2h and the solvent then removed under vacuum and the residue neutralised with saturated K₂CO₃ solution. The aqueous was extracted with EtOAc (200ml) and *n*-butanol (2x200ml) and the combined extracts evaporated under vacuum. The residue was chromatographed on silica gel eluting with CH₂Cl₂/MeOH/NH₃ (60:8:1) to give the title-indole (5.39g, 66%), mp 114-116°C, δ (250MHz, D₆-DMSO) 2.54 (3H, d, J=4.8Hz, MeNH), 2.83 (2H, t, J=7.5Hz, CH₂), 3.60-3.69 (2H, m, CH₂-OH), 4.34 (2H, s, CH₂), 4.65 (1H, t, J=5.4Hz, OH), 6.78 (1H, q, J=4.8Hz, MeNH), 7.06 (1H,

dd, J=2.2 and 8.3Hz, Ar-H), 7.16 (1H, d, J=2.2Hz, Ar-H), 7.31 (1H, d, J=8.3Hz, Ar-H), 7.51 (1H, s, Ar-H), 10.86 (1H, s, NH).

3. (3S)-3-(N-Benzyl)aminomethyl-1-[2-(5-(N-
5 (methyl)aminosulphonylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine 2.1
Hydrogen Oxalate 0.7 Diethyl etherate

Methane sulphonyl chloride (0.32g, 2.80mmol) was added to a stirred solution of 2-[5-(N-(methyl)aminosulphonylmethyl)-1H-indol-3-yl]ethyl alcohol (0.50g, 1.87mmol) and triethylamine (0.38g, 3.73mmol), in
10 dichloromethane (15ml) and acetonitrile (15ml), at 0°C. The mixture was warmed to room temperature and stirred for 3h. Ethyl acetate (70ml) was added to the mixture and the solution washed with water (40ml) and brine (40ml). The organic was dried (MgSO₄) and evaporated. The residue was dissolved in anhydrous acetonitrile (7ml) and IPA (55ml) and
15 Na₂CO₃ (0.26g, 2.5mmol) and Intermediate 1 (0.47g, 2.5mmol) were added. The mixture was heated at reflux for 16h and then cooled to room temperature and the insolubles filtered off. The solvent was removed under vacuum and the residue chromatographed on silica gel eluting with CH₂Cl₂/MeOH/NH₃ (60:8:1) and then again on alumina (Activity III)
20 eluting with CH₂Cl₂/MeOH (98:2) to give the title-ethylpyrrolidine (58mg, 8%). The 2.1 hydrogen oxalate 0.7 diethyl etherate salt was prepared, mp 224-226°C, (Found: C, 54.74, H, 5.99, N, 7.95.

C₂₄H₃₂N₄SO₂·2.1(C₂H₂O₄)·0.7(Et₂O) requires C, 54.63, H, 6.39, N, 8.22%),
m/e 441 (M+1)⁺, δ (360MHz, D₆-DMSO) 1.70-1.84 (1H, m, CH), 2.12-2.26
25 (1H, m, CH), 2.50 (3H, s, MeNH), 2.66-2.80 (1H, m, CH), 2.94-3.56 (10H, m, 5 of CH₂), 4.13 (2H, s, CH₂), 4.34 (2H, s, CH₂), 6.81 (1H, br s, NH), 7.12 (1H, d, J=8.4Hz, Ar-H), 7.26 (1H, s, Ar-H), 7.36 (1H, d, J=8.4Hz, Ar-H), 7.40-7.52 (5H, m, Ar-H), 7.56 (1H, s, Ar-H), 11.04 (1H, s, NH).

EXAMPLE 2**(3S)-3-(N-Benzyl)aminomethyl-1-[2-(5-(aminosulphonylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine 2.0 Hydrogen Oxalate 0.75 Hydrate**

5

1. Intermediate 3: 2-[5-(Aminosulphonylmethyl)-1H-indol-3-yl]ethyl alcohol

Prepared from 4-nitrobenzene methane sulphonamide using the procedures described for Intermediate 2, mp 173-175°C, δ , (D_6 -DMSO)

10 2.83 (2H, t, $J=7.4\text{Hz}$, CH_2), 3.61-3.69 (2H, m, CH_2), 4.29 (2H, s, CH_2SO_2), 4.64 (1H, t, $J=5.3\text{Hz}$, OH), 6.70 (2H, s, NH_2), 7.06 (1H, dd, $J=1.6$ and 8.4Hz , Ar-H), 7.16 (1H, d, $J=1.6\text{Hz}$, Ar-H), 7.31 (1H, d, $J=8.4\text{Hz}$, Ar-H), 7.50 (1H, s, Ar-H), 10.84 (1H, s, NH).

15 **2. (3S)-3-(N-Benzyl)aminomethyl-1-[2-(5-(aminosulphonylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine 2.0 Hydrogen Oxalate 0.75 Hydrate**

Prepared from Intermediates 1 and 3 using the procedure described for Example 1. The 2.0 hydrogen oxalate 0.75 hydrate salt was prepared, mp 203-205°C, (Found: C, 52.52, H, 5.90, N, 8.72;

20 $\text{C}_{23}\text{H}_{30}\text{N}_4\text{SO}_2 \cdot 2(\text{C}_2\text{H}_2\text{O}_4) \cdot 0.75\text{H}_2\text{O}$ requires C, 52.29, H, 5.77, N, 9.03%) m/e 426 ($M+1$)⁺, δ (360MHz, D_6 -DMSO) 1.68-1.82 (1H, m, CH), 2.12-2.24 (1H, m, CH), 2.68-2.78 (1H, m, CH), 2.98-3.54 (10H, m, 5 of CH_2), 4.11 (2H, s, CH_2), 4.31 (2H, s, CH_2), 6.72 (2H, s, NH_2), 7.11 (1H, d, $J=1.5$ and 8.4Hz , Ar-H), 7.25 (1H, d, $J=1.5\text{Hz}$, Ar-H), 7.35 (1H, d, $J=8.4\text{Hz}$, Ar-H), 7.40-7.52
25 (5H, m, Ar-H), 7.55 (1H, s, Ar-H), 11.02 (1H, s, NH).

EXAMPLE 3

30 **(3S)-3-(N-Benzyl)aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine 3.0 Hydrogen Oxalate**

1. Intermediate 4: (S)-2-[5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl alcohol

a) (S)-4-(4-Aminobenzyl)-1,3-oxazolidin-2-one

5 Prepared as described in WO 91/18897.

(b) (S)-4-(3-Iodo-4-aminobenzyl)-1,3-oxazolidin-2-one

A solution of iodine monochloride (4.84g, 29.8mmol) in methanol (35ml) was added dropwise to a stirred mixture of (S)-4-(4-aminobenzyl)-1,3-oxazolidin-2-one (5.2g, 27.0mmol) and calcium carbonate (5.42g, 54.2mmol) in methanol (115ml), at -40°C. The reaction was allowed to warm to room temperature and stir for 16h. The solvent was removed under reduced pressure, the residue taken up into ethyl acetate (300ml) and washed with 20% aqueous sodium thiosulphate (100ml). The organic layer was separated, washed with water (50ml) and brine (50ml), dried (Na₂SO₄) and evaporated. The crude product was chromatographed on silica gel eluting with CH₂Cl₂/MeOH/98:2 to give the title-iodoaniline (3.88, 45%), δ (250MHz, D₆-DMSO) 2.55-2.60 (2H, m, CH₂), 3.90-3.99 (2H, m, CH₂O), 4.19-4.28 (1H, m, CHNH), 5.09 (2H, s, NH₂), 6.69 (1H, d, J=8.2Hz, Ar-H), 6.95 (1H, dd, J=1.9 and 8.2Hz, Ar-H), 7.44 (1H, d, J=1.9Hz, Ar-H). 7.74 (1H, s, NH).

c) (S)-2-[5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl alcohol

25 Prepared from (S)-4-(3-iodo-4-aminobenzyl)-1,3-oxazolidin-2-one and 1,4-bis-triethylsilyl-3-butyn-1-ol as described for Intermediate 2, δ (360MHz, D₆-DMSO) 2.74-2.91 (4H, m, 2 of CH₂), 3.64 (2H, t, J=7.3Hz, CH₂), 4.00-4.08 (2H, m, CH₂), 4.20-4.26 (1H, m, CH), 6.92 (1H, dd, J=1.4 and 8.2Hz, Ar-H), 7.10 (1H, s, Ar-H), 7.25 (1H, d, J=8.2Hz, Ar-H), 7.36 (1H, s, Ar-H), 7.75 (1H, s, NH), 10.69 (1H, s, NH).

2. (3S)-3-(N-Benzyl)aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine. 3.0 Hydrogen Oxalate

Prepared from Intermediates 1 and 4 using the procedure described for Example 1. The 3.0 hydrogen oxalate salt was prepared, mp 196-
5 197°C, (Found: C, 54.46, H, 5.12, N, 7.63. C₂₆H₃₂N₄O₂. 3.0 (C₂H₂O₄) requires C, 54.70, H, 5.45, N, 7.97%), m/e 433 (M+1)⁺, δ (360MHz, D₆-DMSO) 1.72-1.86 (1H, m, CH), 2.14-2.27 (1H, m, CH), 2.68-3.64 (13H, m, 6 of CH₂ and CH), 3.99-4.24 (2H, m, CH₂), 4.16 (2H, s, CH₂), 4.20-4.26 (1H, m, CH), 6.98 (1H, d, J=8.6Hz, Ar-H), 7.20 (1H, s, Ar-H), 7.29 (1H, d,
10 J=8.6Hz, Ar-H), 7.38-7.54 (6H, m, Ar-H), 7.80 (1H, s, NH), 10.91 (1H, s, NH).

EXAMPLE 4

15 (3S)-3-[N-(R)- α -(Hydroxymethyl)benzyl]aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine. 2.2 Hydrogen Oxalate 0.5 Hemihydrate

20 1. Intermediate 5: (3S)-N(H)-3-[α -(Hydroxymethyl)benzyl]aminomethylpyrrolidine

a) (3S)-N-tert-Butyloxycarbonyl-3-[α -(hydroxymethyl)benzyl]aminomethylpyrrolidine

A solution of (R)-(-)-phenylglycinol (2.20g, 16.1mmol) and (3R)-N-
25 *tert*-butyloxycarbonyl-3-methylsulphonyloxymethylpyrrolidine (Intermediate 1 part b; 1.0g, 3.58mmol), in toluene (20ml), was heated at 150°C for 6 h in a sealed pressure tube (Aldrich). The solvent was then removed under vacuum and the residue taken up into ethyl acetate (200ml) and washed with water (x4). The organic was dried (MgSO₄) and
30 evaporated and the crude product chromatographed on silica gel eluting with CH₂Cl₂/MeOH (97:3) to give the title- α -(hydroxymethyl)

benzylaminomethylpyrrolidine (1.0g, 87%), δ (360MHz, CDCl₃) 1.45 (9H, s, OC(Me)₃), 1.52-2.60 (5H, m, CH₂ and CH), 2.90-3.76 (7H, m, 3 of CH₂ and CH), 7.25-7.39 (5H, m, Ar-H).

5 b) (3S)-N(H)-3-[(R)- α -(Hydroxymethyl)benzyl]aminomethylpyrrolidine

Prepared from the preceding N-Boc pyrrolidine using the procedure described for Intermediate 1 part d, δ (250MHz, CDCl₃) 1.25-1.45 (1H, m, CH of CH₂), 1.83-1.97 (1H, m, CH of CH₂), 2.14-2.61 (4H, m, 2 of CH₂), 2.80-3.09 (3H, m, CH₂ and CH), 3.46-3.76 (3H, m, CH₂ and CH), 7.25-7.38 (5H, m, Ar-H).

2. (3S)-3-[N-(R)- α -(Hydroxymethyl)benzyl]aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine 2.2

Hydrogen Oxalate 0.5 Hemihydrate

15 Prepared from Intermediates 4 and 5 using the procedure described for Example 1. The 2.2 hydrogen oxalate 0.5 hemihydrate salt was prepared, mp 115-117°C, (Found: C, 56.37, H, 6.19, N, 8.67. C₂₇H₃₄N₄O₃. 2.2 (C₂H₂O₄). 0.5H₂O requires C, 56.32, H, 5.93, N, 8.37%), m/e 463 (M+1)⁺, δ (360MHz, D₆-DMSO) 1.60-1.76 (1H, m, CH of CH₂), 2.08-2.22 (1H, m, CH of CH₂), 2.46-4.68 (19H, m, 8 of CH₂ and 3 of CH), 6.98 (1H, d, J=8.4Hz, Ar-H), 7.20 (1H, s, Ar-H), 7.29 (1H, d, J=8.4Hz, Ar-H), 7.32-7.46 (6H, m, Ar-H), 7.80 (1H, s, NH), 10.90 (1H, s, NH).

EXAMPLE 5

25 (3S)-3-[N-(S)- α -Methylbenzyl]aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine. 2.4 Hydrogen Oxalate

30 Prepared from (S)-2-[5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl alcohol and (3S)-N(H)-3-(N-(S)- α -methylbenzyl)aminomethyl pyrrolidine using the procedures described for Example 4. The 2.4

hydrogen oxalate salt was prepared, mp 115-117°C, (Found: C, 57.77, H, 5.93, N, 8.77. C₂₇H₃₄N₄O₂ requires C, 57.64, H, 5.90, N, 8.45%), m/e 447 (M+1)⁺. δ (360MHz, D₆-DMSO) 1.51 (2H, d, J=6.7Hz, Me), 1.60-1.72 (1H, m, CH of CH₂), 2.10-2.20 (1H, m, CH of CH₂), 2.48-4.60 (17H, m, 7-CH₂ and 3 of CH), 6.97 (1H, d, J=8.3Hz, Ar-H), 7.19 (1H, s, Ar-H), 7.28 (1H, d, J=8.3Hz, Ar-H), 7.34-7.52 (6H, Ar-H and NH), 7.80 (1H, s, Ar-H), 10.89 (1H, s, NH).

EXAMPLE 6

10

4-[N-(R)- α -(Hydroxymethyl)benzyl]amino-(S)-1-[3-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)propyl]piperidine. 2.15 Hydrogen Oxalate

15

1. (S)-3-[5-(2-oxo-1,3-oxazolidin-4-yl)methyl]-1H-indol-3-yl]propan-1-ol
The title compound was prepared in 61% yield from (S)-4-(3-iodo-4-aminobenzyl)-1,3-oxazolidin-2-one and 1,5-bis-triethylsilyl-4-pentyn-1-ol as described for Intermediate 2. δ (360MHz, DMSO-d₆) 1.78 (2H, gn, J=7.9Hz), 2.69 (2H, t, J=7.4Hz), 2.77 (1H, dd, J=13.5 and 7.1Hz), 2.89 (1H, dd, J=13.5 and 4.6Hz), 3.46 (2H, q, J=5.3Hz), 3.98-4.08 (2H, m), 4.18-4.28 (1H, m), 4.42 (1H, t, J=5.1Hz), 6.92 (1H, dd, J=8.3 and 1.5Hz), 7.06 (1H, d, J=2.1Hz), 7.24 (1H, d, J=8.3Hz), 7.35 (1H, s), 10.66 (1H, s); m/z (ES) 275 (M⁺+1).

20

2. 4-[N-(R)- α -(Hydroxymethyl)benzyl]aminopiperidine

25

To a stirred solution of N-tert-butyloxycarbonyl-4-piperidinone (2g, 10mmol), (R)-(-)-phenylglycinol (1.65g, 12mmol), and glacial acetic acid (2.29ml, 40mmol) in methanol (200ml) was added sodium cyanoborohydride (754mg, 12mmol). After being stirred at room temperature, under nitrogen, for 16 hours, the mixture was basified with 4N sodium hydroxide and the methanol was removed under vacuum. The residue was diluted with water (35ml) and the product extracted with

30

diethyl ether (2x200ml), washed with brine (1x40ml), dried (Na_2SO_4) and concentrated. Flash chromatography (silica gel, dichloromethane-methanol-ammonia, 95:5:0.5) of the residue gave 2.91g (90.9%) of N-tert-butylloxycarbonyl-4-[N-(R)- α -(hydroxymethyl)benzyl]aminopiperidine.

- 5 A solution of the above BOC-protected piperidine (2.9g) in trifluoroacetic acid (40ml) and dichloromethane (50ml) was allowed to stand at room temperature for 16 hours. Solvents were removed under vacuum and the residue was azeotroped with toluene-ethanol (5:1, 150ml). The residue was dissolved in 4N sodium hydroxide, extracted with
10 dichloromethane (3x150ml) and the combined organic solutions were washed with brine (1x50ml), then dried (Na_2SO_4) and concentrated. Crystallisation from ethyl acetate-hexane (1:10, 200ml) afforded the *title compound* as white crystals (1.4g, 70.4%): δ (360MHz, DMSO- d_6) 0.96-1.12 (2H, m), 1.52 (1H, d, $J=12.0$ Hz), 1.78-2.06 (2H, br s and d, $J=12.6$ Hz)
15 2.17-2.32 (3H, m), 2.76-2.90 (2H, m), 3.26 (1H, t, $J=8.5$ Hz), 3.40 (1H, dd, $J=10.5$ and 4.5Hz), 3.83 (1H, dd, $J=8.5$ and 4.5Hz), 4.82 (1H, br s), 7.27-7.37 (5H, m); m/z (ES) 221 (M^++1).

3. 4-[N-(R)- α -(Hydroxymethyl)benzyl]amino-(S)-1-[3-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)propyl]piperidine. 2.15 Hydrogen Oxalate.
20

- The *title compound free base* was prepared from the products of steps 1 and 2 using a similar method to that described for Example 1. The oxalate salt was prepared from ethanol: mp 156-163°C; (Found: C, 57.85; H, 5.97; N, 8.63. $\text{C}_{28}\text{H}_{36}\text{N}_4\text{O}_3 \times 2.15 \text{ C}_2\text{H}_2\text{O}_4$ requires: C, 57.89, H, 6.06; N, 8.36%); m/z (ES) 477 (M^++1); δ (360MHz, DMSO- d_6) 1.66-1.85 (2H, m), 1.92-2.18 (4H, m), 2.62-3.00 (9H, m), 3.30-3.42 (2H, m), 3.58-3.70 (2H, m), 3.98-4.10 (2H, m), 4.14-4.28 (2H, m), 6.95 (1H, d, $J=8.3$ Hz), 7.11 (1H, s), 7.26 (1H, d, $J=8.3$ Hz), 7.30-7.52 (4H, m), 7.79 (1H, s), 10.77 (1H, s).
25
30

EXAMPLE 7

(3S)-3-(N-Benzyl-N-methyl)aminomethyl-(S)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine, Sesquioxalate,

5 Hemihydrate

1. Intermediate 6: (S)-2-[5-(3-Methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl alcohol

10 a) (S)-3-Methyl-4-(4-aminobenzyl)-1,3-oxazolidin-2-one
Prepared as described in WO 91/18897.

b) (S)-2-[5-(3-Methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl alcohol

15 Prepared from (S)-3-methyl-4-(3-iodo-4-aminobenzyl)-1,3-oxazolidin-2-one and 1,4-bis-triethylsilyl-3-butyne-1-ol as described for Intermediate 2, δ (360MHz, D₆-DMSO) 2.72-2.84 (6H, m, CH of CH₂, CH₂ and N-Me), 3.13 (1H, dd, J=3.8 and 13.5Hz, CH of CH₂), 3.61-3.67 (2H, m, CH₂), 3.94-4.02 (2H, m, CH₂), 4.11-4.17 (1H, m, CH), 4.58 (1H, t, J=5.3Hz, OH), 6.93 (1H, dd, J=1.5 and 8.3Hz, Ar-H), 7.10 (1H, d, J=1.5Hz, Ar-H), 7.26 (1H, d, J=8.3Hz, Ar-H), 7.38 (1H, s, Ar-H), 10.72 (1H, s, NH).

25 2. Intermediate 7: (3S)-N(H)-3-(N-Benzyl-N-methyl)-aminomethylpyrrolidine

Prepared from (3R)-N-tert-butyloxycarbonyl-3-hydroxymethylpyrrolidine and N-methylbenzylamine using the procedures described for Intermediate 1, δ (250MHz, D₆-DMSO) 1.41-1.55 (1H, m, CH of CH₂), 1.89-2.02 (1H, m, CH of CH₂), 2.11 (3H, s, Me), 2.31 (2H, d, J=7.5Hz, CH₂NMe), 2.38-2.52 (1H, m, CH), 2.73 (1H, dd, J=6.9 and

11.3Hz, CH of CH₂), 2.95-3.23 (5H, m, 2 of CH₂ and CH of CH₂), 3.46 (2H, ABq, J=13.4Hz, NCH₂-Ar), 7.19-7.36 (5H, m, Ar-H).

3. (3S)-3-(N-Benzyl-N-methyl)aminomethyl-(S)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine.
5 Sesquioxalate. Hemihydrate

Prepared from Intermediates 6 and 7 using the procedure described for Example 1. The sesquioxalate hemihydrate salt was prepared, mp 102-104°C, (Found: C, 61.73; H, 7.02; N, 9.02. C₂₈H₃₆N₄O₂. 1.5(C₂H₂O₄) 0.5H₂O) requires C, 61.58; H, 6.67; N, 9.26%), m/e 461 (M+1)⁺, δ (360MHz, D₆-DMSO) 1.60-1.72 (1H, m, CH of CH₂), 2.08-2.20 (1H, m, CH of CH₂), 2.17 (3H, s, N-Me), 2.44-4.16 (18H, m, 2 of CH and 8 of CH₂), 2.83 (3H, s, N-Me), 7.00 (1H, d, J=8.5Hz, Ar-H), 7.23 (1H, s, Ar-H), 7.24-7.36 (6H, m, Ar-H), 7.44 (1H, s, Ar-H), 10.93 (1H, s, NH).

15

EXAMPLE 8

(3R)-3-[N-(S)-α-Methylbenzyl-N-methylaminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine. 3.0 Hydrogen
20 Oxalate. Hemihydrate

a) (3S)-N-tert-Butyloxycarbonyl-3-[N-(S)-α-methylbenzylaminomethyl
pyrrolidine

Prepared from (3S)-N-tert-butyloxycarbonyl-3-methylsulphonyloxymethylpyrrolidine and (S)-α-methylbenzylamine using the procedure described for Intermediate 5 part a, δ (250MHz, CDCl₃) 1.34 (3H, d, J=6.5Hz, Me), 1.44 (9H, s, OC(Me)₃), 1.44-2.60 (5H, m, 2 of CH₂ and CH), 2.90-3.54 (4H, m, 2 of CH₂), 3.74 (1H, q, J=6.5Hz, CH-Me), 7.18-7.36 (5H, m, Ar-H).

30

b) (3R)-N-tert-Butyloxycarbonyl-3-[N-(S)- α -methylbenzyl-N-methylaminomethylpyrrolidine]

Glacial acetic acid (0.90ml, 15.7mmol) and sodium cyanoborohydride (0.495g, 7.88mmol) were added successively to a stirred solution of the preceding α -(methyl)benzylaminomethylpyrrolidine (1.92g, 6.31mmol) in anhydrous methanol (150ml) at 0°C. A solution of formaldehyde (0.623g, of a 38% w/v solution, 7.88mmol) in methanol (50ml) was then added, dropwise. The mixture was stirred at 0°C for 4.5h and then at room temperature for 1.25h. Saturated K₂CO₃ solution (25ml) was added and the precipitated inorganics were removed by filtration before removing the solvent *in vacuo*. The resultant residue was taken up into ethyl acetate and washed with water (x1) and brine (x2), and dried (MgSO₄). The crude product remaining, after evaporating the solvent *in vacuo*, was chromatographed on silica gel eluting with CH₂Cl₂/MeOH (97.5:2.5) to give the title product (2.02g, 100%), δ (250MHz, CDCl₃) 1.34 (3H, d, J=6.7Hz, Me), 1.44 (9H, s, OC(Me)₃), 1.60-1.68 (1H, m, CH of CH₂), 1.86-1.98 (1H, m, CH of CH₂), 2.19 (3H, s, Me), 2.19-2.42 (3H, m, CH and CH₂), 2.80-3.60 (5H, m, CH and 2 of CH₂), 7.18-7.32 (5H, m, Ar-H).

c) (3R)-N(H)-3-[N-(S)- α -Methylbenzyl-N-methylaminomethylpyrrolidine]

Prepared from the preceding N-Boc pyrrolidine using the procedure described for Intermediate 1, part d, δ (250MHz, CDCl₃) 1.34 (3H, d, J=6.8Hz, Me), 1.52-1.67 (1H, m, CH of CH₂), 1.94-2.08 (1H, m, CH of CH₂), 2.17 (3H, s, Me), 2.20-2.52 (3H, m, CH and CH₂), 2.72 (1H, dd, J=7.3 and 11.3Hz, CH of CH₂), 3.07-3.13 (2H, m, CH₂), 3.25 (1H, dd, J=7.3 and 11.3Hz, CH of CH₂), 3.57 (1H, q, J=6.8Hz, CH-Me), 7.19-7.34 (5H, m, Ar-H).

d) (3R)-3-[N-(S)- α -Methylbenzyl-N-methyl]aminomethyl-(S)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine. 3.0
Hydrogen Oxalate. Hemihydrate.

Triethylamine (0.182ml, 1.3mmol) was added dropwise to a stirred
5 solution of (S)-2-[5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl
alcohol (0.20g, 0.77mmol), in anhydrous THF (12ml). The solution was
cooled to 0°C and methane sulphonyl chloride (0.095ml, 1.2mmol) added
dropwise. The mixture was warmed to room temperature and stirred for
1h before filtering and evaporating the filtrate *in vacuo*. The residue was
10 taken up into dichloromethane (50ml), washed with water (x2) and dried
(MgSO₄). The solvent was removed *in vacuo* to give the desired mesylate
(0.305g) which was used without further purification. Potassium
carbonate (0.159g, 1.15mmol) and sodium iodide (0.115g, 0.767mmol) were
added successively to a stirred solution of the preceding mesylate (0.305g,
15 0.90mmol) in anhydrous DMF (20ml). A solution of (3R)-N-(H)-3-[N-(S)- α -
methylbenzyl-N-methyl]aminomethylpyrrolidine (0.286g, 1.31mmol), in
DMF (5ml), was then added and the mixture heated at 70°C for 18h. The
reaction mixture was cooled to room temperature and then poured into
ethyl acetate (200ml) and washed with water (x6). The organic layer was
20 dried (MgSO₄) and evaporated *in vacuo* to give the crude product which
was chromatographed on silica gel eluting with CH₂Cl₂/MeOH/NH₃
(70:8:1) to give the title-*indole* (59mg, 14%). The 3.0 hydrogen oxalate
hemihydrate salt was prepared, mp 85-90°C (Hygroscopic), (Found: C,
55.26; H, 5.98; N, 7.60. C₂₈H₃₆N₄O₂. 3.0 (C₂H₂O₄). 0.5 H₂O requires C,
25 55.21; H, 5.86; N, 7.57%), m/e 461 (M+1)⁺, δ (360MHz, D₆-DMSO) 1.41 (3H,
d, J=6.8Hz, Me), 1.58-1.70 (1H, m, CH of CH₂), 2.07-2.20 (1H, m, CH of
CH₂), 2.27 (3H, s, Me), 2.50-4.24 (17H, m, 3 of CH and 7 of CH₂), 6.98 (1H,
d, J=8.4Hz, Ar-H), 7.21 (1H, s, Ar-H), 7.26-7.44 (7H, m, Ar-H), 7.80 (1H, s,
Ar-H), 10.90 (1H, s, NH).

EXAMPLE 9

(3R)-3-[N-(S)- α -Methylbenzyl-N-methyl]aminomethyl-(S)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine.

5 2.5 Hydrogen Oxalate. Monohydrate

Prepared from (S)-2-[5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl alcohol (Intermediate 6) and (3R)-N(H)-3-[N-(S)- α -methylbenzyl-N-methyl]aminomethylpyrrolidine using the procedure described for Example 8. The 2.5 hydrogen oxalate monohydrate salt was
10 prepared, low melting point (hygroscopic), (Found: C, 57.01; H, 6.38; N, 7.73. $C_{29}H_{38}N_4O_2 \cdot 2.5(C_2H_2O_4) \cdot 1.0H_2O$ requires C, 56.90; H, 6.32; N, 7.81%), m/e 475 (M+1)⁺, δ (360MHz, D₆-DMSO) 1.42 (3H, d, J=6.8Hz, Me), 1.60-1.74 (1H, m, CH of CH₂), 2.06-2.20 (1H, m, CH of CH₂), 2.27 (3H, s, Me), 2.40-4.20 (17H, m, 3 of CH and 7 of CH₂), 2.84 (3H, s, Me), 7.00 (1H,
15 d, J=8.4Hz, Ar-H), 7.23 (1H, s, Ar-H), 7.32 (1H, d, J=8.4Hz, Ar-H), 7.34-7.40 (5H, m, Ar-H), 7.44 (1H, s, Ar-H), 10.94 (1H, s, NH).

EXAMPLE 10


20 (3S)-3-[N-(4-Fluorobenzyl)-N-methyl]aminomethyl-(S)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl)ethyl]pyrrolidine. 1.65 Hydrogen Oxalate. 0.6 Hydrate

The title-compound was prepared from (S)-2-[5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1H-indol-3-yl]ethyl alcohol and (3S)-3-[N-(4-fluorobenzyl)-N-methyl]aminomethylpyrrolidine using the procedure described for Example 8. The 1.65 hydrogen oxalate 0.6 hydrate salt was
25 prepared, mp 88-89°C, (Found: C, 58.71; H, 6.56; N, 8.82. $C_{28}H_{35}N_4O_2F \cdot 1.65(C_2H_2O_4) \cdot 0.6H_2O$ requires C, 58.93; H, 6.24; N, 8.78%), m/e 479 (M+1)⁺, δ (360MHz, D₆-DMSO) 1.56-1.68 (1H, m, CH of CH₂), 2.04-2.20
30 (4H, m, Me and CH of CH₂), 2.40-4.18 (18H, m, 2 of CH and 8 of CH₂),

2.82 (3H, s, Me), 6.99 (1H, d, J=8.3Hz, Ar-H), 7.12-7.36 (6H, m, Ar-H),
7.43 (1H, s, Ar-H), 10.92 (1H, s, NH).

CLAIMS:

1. A compound of formula I, or a salt or prodrug thereof:




5

(I)


wherein

- Z represents hydrogen, halogen, cyano, nitro, trifluoromethyl, $-OR^5$, $-OCOR^5$, $-OCONR^5R^6$, $-OCH_2CN$, $-OCH_2CONR^5R^6$, $-SR^5$, $-SOR^5$, $-SO_2R^5$, $-SO_2NR^5R^6$, $-NR^5R^6$, $-NR^5COR^6$, $-NR^5CO_2R^6$, $-NR^5SO_2R^6$, $-COR^5$, $-CO_2R^5$, $-CONR^5R^6$, or a group of formula (a), (b), (c) or (d):


10




(a)



(b)



(c)



(d)

- 15 in which the asterisk * denotes a chiral centre;

X represents oxygen, sulphur, $-NH-$ or methylene;

Y represents oxygen or sulphur;

E represents a chemical bond or a straight or branched alkylene chain containing from 1 to 4 carbon atoms;

- 20 Q represents a straight or branched alkylene chain containing from 1 to 4 carbon atoms, optionally substituted in any position by a hydroxy group;

U represents nitrogen or C-R²;

V represents oxygen, sulphur or N-R³;

R², R³ and R⁴ independently represent hydrogen or C₁₋₆ alkyl;

R⁵ and R⁶ independently represent hydrogen, C₁₋₆ alkyl,

5 trifluoromethyl, phenyl, methylphenyl, or an optionally substituted aryl(C₁₋₆)alkyl or heteroaryl(C₁₋₆)alkyl group; or R⁵ and R⁶, when linked through a nitrogen atom, together represent the residue of an optionally substituted azetidine, pyrrolidine, piperidine, morpholine or piperazine ring;

10 M represents the residue of an azetidine, pyrrolidine or piperidine ring;

R represents a group of formula -W-R¹;

W represents a chemical bond or a straight or branched alkylene chain containing from 1 to 4 carbon atoms;

15 R¹ represents -OR^x, -SR^x or -NR^xR^y;


R^x and R^y independently represent hydrogen, hydrocarbon or a heterocyclic group, or R^x and R^y together represent a C₂₋₆ alkylene group; and

20 R^a represents hydrogen, hydroxy, hydrocarbon or a heterocyclic group.

2. A compound as claimed in claim 1 wherein Z represents -SO₂NR⁵R⁶ in which R⁵ and R⁶ are as defined in claim 1.

25 3. A compound as claimed in claim 1 wherein Z represents a group of formula (b) as defined in claim 1.

4. A compound as claimed in claim 1 represented by formula IIA, and salts and prodrugs thereof:



wherein

m is zero, 1, 2 or 3;

5 n is 2, 3 or 4;

p is zero, 1 or 2;


R⁵ and R⁶ are as defined in claim 1;

W¹ represents oxygen, sulphur or N-R¹²; and

10 R¹¹ and R¹² independently represent hydrogen, C₁₋₆ alkyl, aryl, aryl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₆)alkyl, any of which groups may be optionally substituted.

5. A compound as claimed in claim 1 represented by formula IIB, and salts and prodrugs thereof:

15




wherein the asterisk * denotes a chiral centre;

R⁴ and Y are as defined in claim 1; and

20 m, n, p, W¹ and R¹¹ are as defined in claim 4.

6. A compound as claimed in claim 1 represented by formula IIC, and salts and prodrugs thereof:



(IIC)

wherein the asterisk * denotes a chiral centre;

R^4 and Y are as defined in claim 1; and

m, n, p, W^1 and R^{11} are as defined in claim 4.

7. A compound selected from:

(3S)-3-(*N*-benzyl)aminomethyl-1-[2-(5-(*N*-methyl)-aminosulphonylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;

(3S)-3-(*N*-benzyl)aminomethyl-1-[2-(5-(aminosulphonylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;

and salts and prodrugs thereof.

8. A compound selected from:

(3S)-3-(*N*-benzyl)aminomethyl-(*S*)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;

(3S)-3-[*N*-(*R*)- α -(hydroxymethyl)benzyl]aminomethyl-(*S*)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;

(3S)-3-[*N*-(*S*)- α -methylbenzyl]aminomethyl-(*S*)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;

4-[*N*-(*R*)- α -(hydroxymethyl)benzyl]amino-(*S*)-1-[3-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)propyl]piperidine;

and salts and prodrugs thereof.

9. A compound selected from:


- 5 (3*S*)-3-(*N*-benzyl-*N*-methyl)aminomethyl-(*S*)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
(3*R*)-3-[*N*-(*S*)- α -methylbenzyl-*N*-methyl]aminomethyl-(*S*)-1-[2-(5-(2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
(3*R*)-3-[*N*-(*S*)- α -methylbenzyl-*N*-methyl]aminomethyl-(*S*)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
10 (3*S*)-3-[*N*-(4-fluorobenzyl)-*N*-methyl]aminomethyl-(*S*)-1-[2-(5-(3-methyl-2-oxo-1,3-oxazolidin-4-ylmethyl)-1*H*-indol-3-yl)ethyl]pyrrolidine;
and salts and prodrugs thereof.

10. A pharmaceutical composition comprising a compound as
15 claimed in any one of the preceding claims in association with a pharmaceutically acceptable carrier.

11. The use of a compound as claimed in any one of claims 1 to 9
for the manufacture of a medicament for the treatment and/or prevention
20 of clinical conditions for which a subtype-selective agonist of 5-HT_{1D} receptors is indicated.


12. A process for the preparation of a compound as claimed in
any one of claims 1 to 9, which comprises;
25

(A) reacting a compound of formula III:



(III)


wherein Z and E are as defined in claim 1; with a compound of formula IV,
or a carbonyl-protected form thereof:



(IV)


wherein R², Q, M, R and R⁵ are as defined in claim 1; followed, where
required, by N-alkylation by standard methods to introduce the moiety R³;
or

(B) reacting a compound of formula VII:



(VII)

wherein M, R and R⁵ are as defined in claim 1; with a compound of
formula VIII:




(VIII)

wherein Z, E, Q, U and V are as defined in claim 1, and L² represents a suitable leaving group; or

(C) cyclising a compound of formula X:

5




(X)

wherein Z, E, Q, M, R and R' are as defined in claim 1, and D¹ represents a readily displaceable group; followed, where required, by N-alkylation by standard methods to introduce the moiety R³; or

10

(D) cyclising a compound of formula XIII:



(XIII)

15

wherein Z, E, Q, R², M, R and R' are as defined in claim 1, and V¹ represents oxygen or sulphur; and

(E) subsequently, where required, converting a compound of formula I initially obtained into a further compound of formula I by conventional methods.

20

13. A method for the treatment and/or prevention of clinical conditions for which a subtype-selective agonist of 5-HT_{1D} receptors is indicated, which method comprises administering to a patient in need of such treatment an effective amount of a compound as claimed in any one of claims 1 to 9.
- 5

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/GB 95/02759

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D403/06 C07D413/14 A61K31/42 A61K31/405

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GB,A,2 083 463 (GLAXO GROUP LTD) 24 March 1982 see page 1, line 1 - page 2, line 41; claims 1,11 ---	1,10,11, 13
Y	WO,A,91 18897 (WELLCOME FOUND) 12 December 1991 cited in the application see claims ---	1,10,11, 13
Y	WO,A,93 20073 (PFIZER INC) 14 October 1993 see claims 1,15-18 ---	1,10,11, 13
A	WO,A,93 00333 (SMITHKLINE BEECHAM PLC) 7 January 1993 see claims --- -/--	1,10,11, 13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 January 1996

Date of mailing of the international search report

19.0 1.9 6

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340.2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340.3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

Internat. Application No.
PCT/GB 95/02759

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 313 397 (WELLCOME FOUND) 26 April 1989 see claims	1,10,11, 13
A	WO,A,94 02477 (MERCK SHARP & DOHME) 3 February 1994 cited in the application see claims	1,10,11, 13

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB95/02759

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 13 is directed to a method of treatment of the human body
the search has been carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/GB 95/02759

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2083463	24-03-82	AU-B- 550010	27-02-86
		AU-B- 7399581	18-02-82
		BE-A- 889931	11-02-82
		CA-A- 1165765	17-04-84
		CH-A- 652394	15-11-85
		DE-A- 3131752	16-06-82
		FR-A, B 2488606	19-02-82
		JP-B- 1048896	20-10-89
		JP-C- 1563095	12-06-90
		JP-A- 57059865	10-04-82
		NL-A- 8103764	01-03-82
		SE-B- 454777	30-05-88
		SE-A- 8104783	13-02-82
		US-A- 4839377	13-06-89
		AR-A- 231533	28-12-84
		AU-B- 548270	05-12-85
		AU-B- 7399681	18-02-82
		BE-A- 889930	11-02-82
		CA-A- 1169077	12-06-84
		CA-A- 1169429	19-06-84
		CH-A- 651550	30-09-85
		DE-A- 3131748	01-04-82
		FR-A, B 2488605	19-02-82
		GB-A, B 2081717	24-02-82
		JP-B- 1048895	20-10-89
		JP-C- 1565595	25-06-90
		JP-A- 57059864	10-04-82
		LU-A- 83546	08-06-83
		NL-A- 8103768	01-03-82
		SE-B- 454881	06-06-88
		SE-A- 8104782	13-02-82
		US-A- 4672067	09-06-87
		US-A- 4636521	13-01-87
WO-A-9118897	12-12-91	AU-B- 646871	10-03-94
		AU-B- 7957091	31-12-91
		CA-A- 2064815	08-12-91
		EP-A- 0486666	27-05-92
		EP-A- 0636623	01-02-95
		JP-T- 5502679	13-05-93

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/LB 95/02759

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9118897		LT-A, B 419	25-11-94
		LV-B- 10274	20-04-95
		NZ-A- 238424	23-12-93
		PL-B- 166214	28-04-95
		US-A- 5466699	14-11-95
		US-A- 5399574	21-03-95
WO-A-9320073	14-10-93	AU-B- 3789893	08-11-93
		CA-A- 2133413	14-10-93
		CN-A- 1082543	23-02-94
		CZ-A- 9402464	14-06-95
		EP-A- 0635017	25-01-95
		HR-A- 930752	31-10-94
		JP-T- 7504919	01-06-95
		NO-A- 943762	06-10-94
		SK-A- 120694	10-05-95
WO-A-9300333	07-01-93	AU-B- 1928892	25-01-93
		CA-A- 2110574	07-01-93
		EP-A- 0593513	27-04-94
		JP-T- 6508354	22-09-94
		ZA-A- 9204523	20-12-93
EP-A-0313397	26-04-89	AU-B- 2418188	27-04-89
		DE-A- 3881472	08-07-93
		DE-T- 3881472	16-12-93
		ES-T- 2054825	16-08-94
		JP-A- 1146882	08-06-89
		US-A- 5225431	06-07-93
WO-A-9402477	03-02-94	AU-B- 4578593	14-02-94
		CA-A- 2138649	03-02-94
		EP-A- 0651749	10-05-95
		JP-T- 7509452	19-10-95